

# Recombinant Rat Cd40 protein, C-His Tag

## Product Information

---

<b>Cat</b>	IMP-376
<b>Official Symbol</b>	Cd40
<b>Product Overview</b>	Recombinant Rat Cd40 protein (NP_599187.1) (Met1-Arg193) was expressed in HEK293, fused with a polyhistidine tag at the C-terminus.
<b>Description</b>	CD40, also known as TNFRSF5, is a member of the TNF receptor superfamily which are single transmembrane-spanning glycoproteins. CD40 protein plays an essential role in mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. CD40 protein is expressed in B cells, dendritic cells, macrophages, endothelial cells, and several tumor cell lines. Defects in CD40 result in hyper-IgM immunodeficiency type 3 (HIGM3). In addition, CD40/CD40L interaction is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis.
<b>Expression System</b>	HEK293
<b>Species</b>	Rat
<b>Tag</b>	C-His Tag
<b>Predicted N Terminal</b>	Leu 20
<b>Form</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
<b>Molecular Mass</b>	The recombinant rat CD40 comprises 185 amino acids and predicts a molecular mass of 20.6 kDa. The apparent molecular mass of the recombinant protein is approximately 28 kDa in SDS-PAGE under reducing conditions due to glycosylation.
<b>Protein length</b>	Met1-Arg193
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method
<b>Purity</b>	> 95 % as determined by SDS-PAGE
<b>Storage</b>	Samples are stable for up to twelve months from date of receipt at -20°C to -80°C. Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover

*the entire contents.*