



Recombinant Human LAG3 Protein, No Tag

Product Information

Cat	IMP-4788
Official Symbol	LAG3
Product Overview	Recombinant human LAG3 (P18627) (Met1-Arg440) was expressed with an initial Met.
Description	LAG3 (Lymphocyte Activating 3) is a Protein Coding gene. 2 alternatively spliced human isoforms have been reported. LAG3, also known as CD223 and Lymphocyte activation gene 3, belongs to the immunoglobulin (Ig) superfamily. The LAG3 gene contains 8 exons. It is selectively expressed in activated T and NK cells. LAG3 contains 4 extracellular Ig-like domains and has a negative regulatory function in T cells. It also acts as a new marker of T cell-induced B cell activation. As a soluble molecule, LAG3 activates antigen-presenting cells through MHC class II signaling, leading to increased antigen-specific T-cell responses in vivo. Diseases associated with LAG3 include Smoldering Myeloma and Kyphoscoliotic Heart Disease.
Expression System	HEK293
Species	Human
Tag	No Tag
Predicted N Terminal	Leu 23
Form	Lyophilized from sterile PBS, pH 7.4, 5 % trehalose, 5% mannitol and 0.01% Tween80.
Molecular Mass	The recombinant human LAG3 consists 418 amino acids and predicts a molecular mass of 45.3 kDa.
Protein length	Met1-Arg440
Endotoxin	< 1.0 EU/μg of the protein as determined by the LAL method
Purity	> 90 % as determined by SDS-PAGE. > 95 % as determined by SEC-HPLC.
Storage	Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade. Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
SDS-PAGE	



HPLC



97.2% as determined by SEC-HPLC Analysis