

Recombinant Mouse Lep Protein, C-Myc/DDK-tagged

Product Information

Cat	IMP-6905
Official Symbol	LEP
Product Overview	Purified recombinant protein of Mouse leptin (Lep), with C-terminal Myc/DDK tag, expressed in HEK293T cells.
Description	 Key player in the regulation of energy balance and body weight control. Once released into the circulation, has central and peripheral effects by binding LEPR, found in many tissues, which results in the activation of several major signaling pathways (PubMed: 15899045, PubMed: 16825198, PubMed: 11373681, PubMed: 12594516, PubMed: 20620997). In the hypothalamus, acts as an appetite-regulating factor that induces a decrease in food intake and an increase in energy consumption by inducin anorexinogenic factors and suppressing orexigenic neuropeptides, also regulates bone mass and secretion of hypothalamo-pituitary-adrenal hormones. In the periphery, increases basal metabolism, influences reproductive function, regulates pancreatic beta-cell function and insulin secretion, is pro-angiogenic for endothelial cell and affects innate and adaptive immunity (By similarity) (PubMed: 8589726, PubMed: 10660043, PubMed: 25383904, PubMed: 25060689, PubMed: 9732873, PubMed: 12594516). In the arcuate nucleus of the hypothalamus, activates by depolarization POMC neurons inducing FOS and SOCS3 expression to release anorexigenic peptides and inhibits by hyperpolarization NPY neurons inducing SOCS3 with a consequent reduction on release of orexigenic peptides (By similarity) (PubMed: 20620997, PubMed: 11373681). In addition to its known satiety inducing effect, has a modulatory role in nutrient absorption. In the intestine, reduces glucose absorption by enterocytes by activating PKC and leading to a sequential activation of p38, PI3K and ERK signaling pathways increases expression of genes involved in cell cycle regulation such as CCND1, via JAK2-STAT3 pathway, or VEGFA, via MAPK1/3 and PI3K-AKT1 pathways (By similarity) (PubMed: 16825198, PubMed: 20620997). May also play ar apoptotic role via JAK2-STAT3 pathway and up-regulation of BIRC5 expression (By similarity). Pro-angiogenic, has mitogenic activity on vascular endothelial cells and plays a role in matrix remodeling by regulating the expression of matrix metalloprotei



	JAK2, PI3K, MAP2K1/MEK1 and MAPK14/p38 (PubMed: 15899045). In adaptive immunity, promotes the switch of memory T-cells towards T helper-1 cell immune responses (By similarity). Increases CD4(+)CD25(-) T cells proliferation and reduces autophagy during TCR (T cell receptor) stimulation, through MTOR signaling pathway activation and BCL2 up- regulation (PubMed: 25060689).[UniProtKB/Swiss-Prot Function]
Expression System	HEK293T
Species	Mouse
Tag	C-Myc/DDK
Form	25mM Tris.HCl, pH 7.3, 100mM glycine, 10% glycerol
Molecular Mass	18.7 kDa
AA Sequence	MCWRPLCRFLWLWSYLSYVQAVPIQKVQDDTKTLIKTIVTRINDISHTQSV SAKQRVTGLDFIPGLHPILSLSKMDQTLAVYQQVLTSLPSQNVLQIANDLE NLRDLLHLLAFSKSCSLPQTSGLQKPESLDGVLEASLYSTEVVALSRLQG SLQDILQQLDVSPECTRTRPLEQKLISEEDLAANDILDYKDDDDKV
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining.
Storage	Store at -80 centigrade after receiving vials. Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
Concentration	> 50 μ g/mL as determined by microplate BCA method.